



Target Communities and Condition for Forest Land Reclamation

(2011)

Kevin Kemball¹

JP Bielich¹

¹ Incremental Forest Technologies



Table of Contents

1. Introduction	4
2. Selecting a Target.....	4
2.1. Target Communities of the Boreal Mixedwood.....	7
Lichen	7
Blueberry	7
Labrador tea–mesic	8
Low-bush cranberry	8
Dogwood.....	9
Horsetail	10
Labrador tea–subhygric	10
Labrador tea/horsetail	11
Bog	11
Poor fen	11
Rich fen	12
2.2. Target Communities of the Lower Foothills	12
Grassland	12
Bearberry/lichen.....	13
Hairy wild rye	13
Labrador tea–mesic	14
Low-bush cranberry	14
Bracted honeysuckle.....	15
Meadow	16
Labrador tea–subhygric	16
Horsetail	17
Labrador tea/horsetail	17
Bog	18
Poor fen	18



Target Plant Communities

Rich fen	19
Marsh.....	19
3. Autecology of key Boreal Mixedwood species	19
Adapted from <i>Alberta Silviculture Guide</i> (2007)	19



1. Introduction

The target community or condition is the expected end point of reclamation. Selection of a target community incorporates the core concept of equivalent land capability. The Conservation and Reclamation Regulation defines equivalent land capability where “the ability of the land to support various land uses after conservation and reclamation is similar to the ability that existed prior to an activity being conducted on the land, but that the individual land uses will not necessarily be identical.”

As set out in the 2010 Reclamation Criteria for Forested Lands, selection of a target community or condition that is different from the adjacent undisturbed land would be a non-routine application. Non-routine applications have modified sampling and reporting criteria and must be justified based on site conditions or other identified constraints. Non-routine applications will also receive additional scrutiny from regulators.

Once selected the target community will act as a filter that will require certain actions be taken, and constrain other reclamation activities that may limit the establishment and growth of the desired plants (particularly woody plants). Based on the initial assessment, some sites may require that the target be different than the pre-disturbance condition (it is non-routine). The most likely change in condition is from lowland to upland (wet site to dryer site), or from a poor to a richer condition. Change in condition is most often dictated by the type and amount of fill material (clay or coarse aggregate material used for pad or road construction) remaining on site.

Note: condition refers to the underlying site capability as described by moisture and nutrient status.

2. Selecting a Target

The selection of a target community for reclamation is the most important step in planning and flows from site assessment. Central to selecting a target for reclamation is understanding how the two main components of the 2010 Reclamation Criteria will be implemented. The two main components are:

1. Assessment Process and Criteria: This outlines the standards and methodology used for assessing various components of the Landscape, Vegetation, and Soil Assessments that make up the 2010 Reclamation Criteria.
2. Assessment Tool and Record of Observation (RoO) Datasheets: The Assessment Tool for each land type poses Yes/No questions based on what is present onsite compared to offsite and whether it meets the standard set out in the Rationale.

Combined, the assessment process and the RoO are used to support the application for a reclamation certificate and to identify areas of noncompliance.

The first step in selecting a target community is assessing the site within the context of the surrounding landscape. Guidance for the required landscape assessment is provided in the



Target Plant Communities

2010 Reclamation Criteria for Forested Lands. The required landscape level assessment includes:

1. Drainage
 - a. Surface Water Flow;
 - b. Sub-Surface Water Flow;
 - c. Riparian Areas
2. Erosion
3. Stability (Subsidence)
4. Bare Areas
5. Operability
 - a. Macro-contour;
 - b. Meso-contour;
 - c. Micro-contour;
 - d. Surface Stoniness
6. Debris
 - a. Woody Debris;
 - b. Refuse

When assessing the landscape prior to target community selection the goal is to identify potential future problems and opportunities. Depending on the age and type of disturbance the site may be partially re-vegetated. Some sites may have topsoil placement complete and others have no soil work conducted. Regardless of the stage of abandonment, all six required criteria for the landscape assessment must be completed. If site adjustment is required based on the landscape assessment then the assessment must be repeated after work is complete. Whenever possible it is highly recommended that the landscape assessment be conducted after sufficient time has passed since site adjustment (e.g., drainage repair or modification, soil placement) for potential landscape issues to manifest.

The next step in selecting a target community is to assess soil and vegetation. Soil assessment must be performed both on site and at reference (control) sites. Depending on the age and stage of abandonment, vegetation assessments may also include both on and off site data collection. The intensity, location and timing of data collection should be similar to that described in the 2010 Reclamation Criteria for Forested Lands. Use of the standards and methodology described in the 2010 Reclamation Criteria will allow for early identification of noncompliance. Early identification will allow for additional remediation or site adjustment to occur and avoid costly duplication of other reclamation work. During the assessment, emphasis should be placed on determining the ecosite and ecosite phase of the surrounding undisturbed lands using the *Field Guide to Ecosites of Northern Alberta* (Beckingham and Archibold 1996). In addition, in the surrounding undisturbed landscape, the presence, age and condition of trees and shrubs that may provide seed should be noted. This will allow natural regeneration to be integrated into the reclamation plan.



Target Plant Communities

If the condition of the site has changed substantially compared to the immediately surrounding landscape, it may not be necessary to sample adjacent lands. In this case placement of sample plots may be expanded to include stands further away but likely more representative of the target community. For example, if a proposed reclamation site was formerly a peatland and is now an upland site, then sampling can be conducted in undisturbed upland sites farther away from the reclamation site. Placement of additional sampling locations along the access route to the site is recommended.

Once the landscape, soil and vegetation data have been compiled, selecting the target community is possible. When no landscape or soil issues are found, selecting a target community can be done in the field by experienced personnel. Selecting the target community should consider the following questions in order. Answering yes to any of the questions below would require that additional sampling be conducted prior to selecting a target plant community, or that a non-routine application will be planned:

1. Are there any landscape issues that must be corrected?
2. Were any of the required soil criteria in noncompliance?
3. Will the site no longer support the adjacent community (condition change)?
4. Would the site better support a community other than the adjacent community?
5. Is there native vegetation on the site that is established, healthy and providing some measure of restored ecosystem function (vegetation override, or minor adjustment)?

The starting point for target community selection is the plant community present on adjacent undisturbed land as described by ecosite and ecosite phase (Beckingham and Archibold 1996). When selecting a target community, the first decision must be based on ecosite. Ecosite is determined from the position on the edatope (moisture regime and nutrient regime). Position on the edatope must be estimated using information collected during the assessment. Most ecosites will support more than one tree species and multiple shrub species. For example, in the boreal mixedwood (*Field Guide to Ecosites of Northern Alberta*), only ecosite a (lichen) and ecosite k (rich fen) contain only one characteristic tree species. All other ecosites support multiple tree species that may occur together in highly variable mixtures. Within an ecosite, the understory communities are typically composed of the same suite of species that differ primarily in abundance. Ecosite phase is based on the tallest vegetation layer (typically trees) and dominant lower vegetation. It is not recommended that the plant community type (the finest level of resolution in the Alberta ecosite classification system) be used for target selection. Plant community types are highly variable and descriptive of large homogeneous mature forest stands.

If a site will not support the trees and shrubs of the surrounding undisturbed land, then selection of the target community should be based on the assessed site conditions, and on anticipated future condition. Future conditions are largely driven by landscape position and surrounding terrain (e.g., position in drainage basin, position on slope).



2.1. Target Communities of the Boreal Mixedwood

The following summarizes the woody vegetation characteristic of ecosites and ecosite phases in the Boreal Mixedwood of northern Alberta, as explained in detail in *Field Guide to Ecosites of Northern Alberta* (Beckingham and Archibald 1996). These summaries, and the accompanying tables, are intended to help set targets for reclaimed sites based on moisture and nutrient regimes, as well as vegetation and wildlife objectives.

Lichen

This ecosite is characterized by xeric to subxeric moisture regimes and poor to very poor nutrient levels. Indicator species include jack pine, blueberry, bearberry, bog cranberry, sand heather and lichen. This ecosite has one ecosite phase, (a1) lichen Pj.

(a) Lichen – characteristic woody plant species.

Tree	<i>Pinus banksiana</i> (jack pine)
Shrub	<i>Arctostaphylos uva-ursi</i> (bearberry) <i>Vaccinium myrtilloides</i> (blueberry) <i>Alnus crispa</i> (green alder) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Ledum groenlandicum</i> (Labrador tea) <i>Linnaea borealis</i> (twin-flower) <i>Pinus banksiana</i> (jack pine) <i>Hudsonia tomentosa</i> (sand heather)

Blueberry

This ecosite is characterized by submesic to subxeric moisture regimes and medium to poor nutrient levels. Indicator species include blueberry, bearberry, bog cranberry, Labrador tea, cream-colored vetchling and hairy wild rye. This ecosite has four ecosite phases, (b1) blueberry Pj-Aw, (b2) blueberry Aw(Bw), (b3) blueberry Aw-Sw, and (b4) blueberry Sw-Pj.

(b) Blueberry – characteristic woody plant species.

Tree	<i>Pinus banksiana</i> (jack pine) <i>Populus tremuloides</i> (aspen) <i>Betula papyrifera</i> (white birch) <i>Picea glauca</i> (white spruce) <i>Picea mariana</i> (black spruce)
Shrub	<i>Ledum groenlandicum</i> (Labrador tea) <i>Alnus crispa</i> (green alder) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Vaccinium myrtilloides</i> (blueberry) <i>Arctostaphylos uva-ursi</i> (bearberry) <i>Picea glauca</i> (white spruce) <i>Shepherdia canadensis</i> (Canada buffalo-berry)



Linnaea borealis (twin-flower)
Populus tremuloides (aspen)
Rosa acicularis (prickly rose)
Picea mariana (black spruce)
Amelanchier alnifolia (saskatoon)
Salix spp. (willow)
Betula papyrifera (white birch)

Labrador tea–mesic

This ecosite is characterized by subhygric to submesic moisture regimes and poor to very poor nutrient levels and relatively acidic surface soil conditions. Indicator species include jack pine, black spruce, Labrador tea, bog cranberry, blueberry, feather moss and reindeer lichen. This ecosite has one ecosite phase, (c1) Labrador tea–mesic Pj-Sb.

(c) Labrador tea–mesic– characteristic woody plant species.

Tree	<i>Pinus banksiana</i> (jack pine)
	<i>Picea mariana</i> (black spruce)
Shrub	<i>Ledum groenlandicum</i> (Labrador tea)
	<i>Alnus crispa</i> (green alder)
	<i>Vaccinium vitis-idaea</i> (bog cranberry)
	<i>Vaccinium myrtilloides</i> (blueberry)
	<i>Linnaea borealis</i> (twin-flower)
	<i>Picea mariana</i> (black spruce)

Low-bush cranberry

This ecosite is characterized by a mesic moisture regime and medium nutrient levels. Indicator species include low-bush cranberry, Canada buffalo-berry, dewberry and wild sarsaparilla. This ecosite has three ecosite phases, (d1) low-bush cranberry Aw, (d2) low-bush cranberry Aw-Sw, and (d3) low-bush cranberry Sw.

(d) Low-bush cranberry – characteristic woody plant species.

Tree	<i>Populus tremuloides</i> (aspen)
	<i>Populus balsamifera</i> (balsam poplar)
	<i>Betula papyrifera</i> (white birch)
	<i>Abies balsamea</i> (balsam fir)
	<i>Picea glauca</i> (white spruce)
	<i>Picea mariana</i> (black spruce)
Shrub	<i>Corylus cornuta</i> (beaked hazelnut)
	<i>Alnus crispa</i> (green alder)
	<i>Rosa acicularis</i> (prickly rose)
	<i>Shepherdia canadensis</i> (Canada buffalo-berry)
	<i>Amelanchier alnifolia</i> (saskatoon)



Abies balsamea (balsam fir)
Viburnum edule (low-bush cranberry)
Salix spp. (willow)
Linnaea borealis (twin-flower)
Populus tremuloides (aspen)
Prunus spp. (pin and choke cherry)
Rubus idaeus (wild red raspberry)
Symphoricarpos albus (snowberry)
Picea glauca (white spruce)
Ribes spp. (currant)
Picea mariana (black spruce)
Betula papyrifera (white birch)
Populus balsamifera (balsam poplar)

Dogwood

This ecosite is characterized by a subhygric moisture regime and rich nutrient levels. Indicator species include balsam poplar, bracted honeysuckle, dogwood, currants, wild red raspberry, tall lungwort, sweet-scented bedstraw, and oak, lady and shield ferns. This ecosite has three ecosite phases, (e1) dogwood Pb-Aw, (e2) dogwood Pb-Sw, and (e3) dogwood Sw. The dogwood ecosite tends to be the most productive in the Boreal Mixedwood.

(e) Dogwood – characteristic woody plant species.

Tree	<i>Populus tremuloides</i> (aspen) <i>Populus balsamifera</i> (balsam poplar) <i>Betula papyrifera</i> (white birch) <i>Picea glauca</i> (white spruce) <i>Abies balsamea</i> (balsam fir)
Shrub	<i>Alnus tenuifolia</i> (river alder) <i>Cornus stolonifera</i> (dogwood) <i>Lonicera involucrata</i> (bracted honeysuckle) <i>Rosa acicularis</i> (prickly rose) <i>Viburnum edule</i> (low-bush cranberry) <i>Salix</i> spp. (willow) <i>Alnus crispa</i> (green alder) <i>Amelanchier alnifolia</i> (saskatoon) <i>Ribes</i> spp. (currant) <i>Linnaea borealis</i> (twin-flower) <i>Populus balsamifera</i> (balsam poplar) <i>Rubus idaeus</i> (wild red raspberry) <i>Picea glauca</i> (white spruce) <i>Abies balsamea</i> (balsam fir) <i>Betula papyrifera</i> (white birch) <i>Alnus crispa</i> (green alder) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Ledum groenlandicum</i> (Labrador tea)



Populus tremuloides (aspen)

Horsetail

This ecosite is characterized by hygric to subhygric moisture regimes and rich to very rich nutrient levels. Indicator species include common horsetail and meadow horsetail. This ecosite has three ecosite phases, (f1) horsetail Pb-Aw, (f2) horsetail Pb-Sw, and (f3) horsetail Sw.

(f) Horsetail – characteristic woody plant species.

Tree	<i>Populus tremuloides</i> (aspen) <i>Populus balsamifera</i> (balsam poplar) <i>Betula papyrifera</i> (white birch) <i>Picea glauca</i> (white spruce) <i>Abies balsamea</i> (balsam fir)
Shrub	<i>Salix</i> spp. (willow) <i>Rosa acicularis</i> (prickly rose) <i>Alnus</i> spp. (green and river alder) <i>Cornus stolonifera</i> (dogwood) <i>Rubus idaeus</i> (wild red raspberry) <i>Viburnum edule</i> (low-bush cranberry) <i>Populus balsamifera</i> (balsam poplar) <i>Picea glauca</i> (white spruce) <i>Abies balsamea</i> (balsam fir) <i>Linnaea borealis</i> (twin-flower) <i>Betula papyrifera</i> (white birch) <i>Ribes</i> spp. (currant) <i>Populus tremuloides</i> (aspen)

Labrador tea–subhygric

This ecosite is characterized by hygric to subhygric moisture regimes and medium to very poor nutrient levels. Indicator species include black spruce, jack pine, Labrador tea and bog cranberry. This ecosite has one ecosite phase, (g1) Labrador tea–subhygric Sb-Pj.

(g) Labrador tea–subhygric – characteristic woody plant species.

Tree	<i>Picea mariana</i> (black spruce) <i>Pinus banksiana</i> (jack pine)
Shrub	<i>Ledum groenlandicum</i> (Labrador tea) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Picea mariana</i> (black spruce) <i>Vaccinium myrtilloides</i> (blueberry) <i>Rosa acicularis</i> (prickly rose) <i>Linnaea borealis</i> (twin-flower)



Labrador tea/horsetail

This ecosite is characterized by hygric to subhygric moisture regimes and medium to rich nutrient levels. Indicator species include white spruce, black spruce, Labrador tea, common horsetail, meadow horsetail and woodland horsetail. This ecosite has one ecosite phase, (h1) Labrador tea/horsetail Sw-Sb.

(h) Labrador tea/horsetail – characteristic woody plant species.

Tree	<i>Picea glauca</i> (white spruce) <i>Picea mariana</i> (black spruce) <i>Betula papyrifera</i> (white birch)
Shrub	<i>Ledum groenlandicum</i> (Labrador tea) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Salix</i> spp. (willow) <i>Rosa acicularis</i> (prickly rose) <i>Picea mariana</i> (black spruce) <i>Linnaea borealis</i> (twin-flower) <i>Populus tremuloides</i> (aspen) <i>Betula papyrifera</i> (white birch)

Bog

This ecosite is characterized by hydric to subhydric moisture regimes and poor to very poor nutrient levels. Indicator species include black spruce, Labrador tea, bog cranberry, cloudberry and peat moss. This ecosite has two ecosite phases, (i1) treed bog and (i2) shrubby bog.

(i) Bog – characteristic woody plant species.

Tree	<i>Picea mariana</i> (black spruce)
Shrub	<i>Ledum groenlandicum</i> (Labrador tea) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Picea mariana</i> (black spruce) <i>Oxycoccus microcarpus</i> (small bog cranberry) <i>Chamaedaphne calyculata</i> (leatherleaf)

Poor fen

This ecosite is characterized by hydric to subhydric moisture regimes and medium to very poor nutrient levels. Indicator species include black spruce, tamarack, Labrador tea, dwarf birch, willow, cloudberry, sedge, peat moss, golden moss, tufted moss and brown moss. This ecosite has two ecosite phases, (j1) treed poor fen and (j2) shrubby poor fen.

(j) Poor fen – characteristic woody plant species.

Tree	<i>Picea mariana</i> (black spruce) <i>Larix laricina</i> (tamarack)
------	---



Shrub	<i>Ledum groenlandicum</i> (Labrador tea) <i>Picea mariana</i> (black spruce) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Salix</i> spp. (willow) <i>Larix laricina</i> (tamarack) <i>Betula pumila</i> (dwarf birch) <i>Oxycoccus microcarpus</i> (small bog cranberry)
-------	--

Rich fen

This ecosite is characterized by hydric to hygric moisture regimes and medium to very rich nutrient levels. Indicator species include tamarack, willow, dwarf birch, sedge, golden moss, tufted moss and brown moss. This ecosite has three ecosite phases, (k1) treed rich fen, (k2) shrubby rich fen and (k3) graminoid rich fen. There are no characteristic woody plant species found in (k3) graminoid rich fen.

(k) Rich fen – characteristic woody plant species (k1 and k2 only).

Tree	<i>Larix laricina</i> (tamarack) <i>Picea mariana</i> (black spruce)
Shrub	<i>Betula pumila</i> (dwarf birch) <i>Larix laricina</i> (tamarack) <i>Salix</i> spp. (willow) <i>Ledum groenlandicum</i> (Labrador tea) <i>Andromeda polifolia</i> (bog rosemary) <i>Picea mariana</i> (black spruce) <i>Alnus tenuifolia</i> (river alder) <i>Cornus stolonifera</i> (dogwood) <i>Ribes</i> spp. (currant) <i>Rubus idaeus</i> (wild red raspberry)

2.2. Target Communities of the Lower Foothills

The following section summarizes the ecosites and ecosite phases in the Lower Foothills of northwestern Alberta, as explained in detail in *Field Guide to Ecosites of West-central Alberta* (Beckingham et al. 1996). These summaries, and the accompanying tables, are intended to help set targets for reclaimed sites based on moisture and nutrient regimes, as well as vegetation and wildlife objectives.

Grassland

This ecosite is characterized by xeric to subxeric moisture regimes and medium to very poor nutrient levels. Indicator species include bearberry, juniper, slender wheat grass and hairy wild rye. This ecosite has one ecosite phase, (a1) shrubby grassland.

**(a) Grassland – characteristic woody plant species.**

Tree	No tree species
Shrub	<i>Arctostaphylos uva-ursi</i> (bearberry) <i>Shepherdia canadensis</i> (Canada buffalo-berry) <i>Juniperus</i> spp. (juniper) <i>Potentilla fruticosa</i> (shrubby cinquefoil) <i>Amelanchier alnifolia</i> (saskatoon) <i>Populus tremuloides</i> (aspen) <i>Rosa acicularis</i> (prickly rose)

Bearberry/lichen

This ecosite is characterized by submesic to xeric moisture regimes and poor to very poor nutrient levels. Indicator species include lodgepole pine, bearberry, blueberry, bog cranberry and lichen. This ecosite has one ecosite phase, (b1) bearberry/lichen PI.

(b) Bearberry/lichen – characteristic woody plant species.

Tree	<i>Pinus contorta</i> (lodgepole pine) <i>Picea mariana</i> (black spruce) <i>Picea glauca</i> (white spruce)
Shrub	<i>Arctostaphylos uva-ursi</i> (bearberry) <i>Rosa acicularis</i> (prickly rose) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Vaccinium myrtilloides</i> (blueberry) <i>Picea mariana</i> (black spruce) <i>Linnaea borealis</i> (twin-flower) <i>Vaccinium caespitosum</i> (dwarf bilberry) <i>Shepherdia canadensis</i> (Canada buffalo-berry) <i>Salix</i> spp. (willow) <i>Ledum groenlandicum</i> (Labrador tea)

Hairy wild rye

This ecosite is characterized by submesic to subxeric moisture regimes and medium to poor nutrient levels. Indicator species include Canada buffalo-berry, blueberry, bearberry, Labrador tea, and hairy wild rye. This ecosite has four ecosite phases, (c1) hairy wild rye PI, (c2) hairy wild rye Aw, (c3) hairy wild rye Aw-Sw-PI, and (c4) hairy wild rye Sw.

(c) Hairy wild rye– characteristic woody plant species.

Tree	<i>Pinus contorta</i> (lodgepole pine) <i>Populus tremuloides</i> (aspen) <i>Picea glauca</i> (white spruce) <i>Populus balsamifera</i> (balsam poplar)
------	--



	<i>Betula papyrifera</i> (white birch)
Shrub	<i>Alnus crispa</i> (green alder) <i>Shepherdia canadensis</i> (Canada buffalo-berry) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Linnaea borealis</i> (twin-flower) <i>Arctostaphylos uva-ursi</i> (bearberry) <i>Rosa acicularis</i> (prickly rose) <i>Ledum groenlandicum</i> (Labrador tea) <i>Picea mariana</i> (black spruce) <i>Vaccinium myrtilloides</i> (blueberry) <i>Populus tremuloides</i> (aspen) <i>Symphoricarpos albus</i> (snowberry) <i>Salix</i> spp. (willow) <i>Lonicera dioica</i> (twining honeysuckle) <i>Viburnum edule</i> (low-bush cranberry)

Labrador tea–mesic

This ecosite is characterized by subhygric to submesic moisture regimes and poor to very poor nutrient levels. Indicator species include lodgepole pine, black spruce, Labrador tea, bog cranberry and blueberry. This ecosite has one ecosite phase, (d1) Labrador tea–mesic PI-Sb.

(d) Labrador tea–mesic– characteristic woody plant species.

Tree	<i>Pinus contorta</i> (lodgepole pine) <i>Picea mariana</i> (black spruce)
Shrub	<i>Ledum groenlandicum</i> (Labrador tea) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Alnus crispa</i> (green alder) <i>Vaccinium myrtilloides</i> (blueberry) <i>Linnaea borealis</i> (twin-flower) <i>Picea mariana</i> (black spruce)

Low-bush cranberry

This ecosite is characterized by subhygric to submesic moisture regimes and rich to poor nutrient levels. Indicator species include low-bush cranberry, dewberry and wild sarsaparilla. This ecosite has four ecosite phases, (e1) low-bush cranberry PI, (e2) low-bush cranberry Aw, (e3) low-bush cranberry Aw-Sw-PI, and (e4) low-bush cranberry Sw.

(e) Low-bush cranberry – characteristic woody plant species.

Tree	<i>Pinus contorta</i> (lodgepole pine) <i>Picea glauca</i> (white spruce) <i>Populus tremuloides</i> (aspen) <i>Populus balsamifera</i> (balsam poplar)
------	--



	<i>Betula papyrifera</i> (white birch) <i>Pinus contorta</i> (lodgepole pine) <i>Abies balsamea</i> (balsam fir) <i>Abies lasiocarpa</i> (subalpine fir) <i>Picea mariana</i> (black spruce)
Shrub	<i>Alnus crispa</i> (green alder) <i>Viburnum edule</i> (low-bush cranberry) <i>Linnaea borealis</i> (twin-flower) <i>Rosa acicularis</i> (prickly rose) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Picea glauca</i> (white spruce) <i>Amelanchier alnifolia</i> (saskatoon) <i>Shepherdia canadensis</i> (Canada buffalo-berry) <i>Salix</i> spp. (willow) <i>Rubus idaeus</i> (wild red raspberry) <i>Populus tremuloides</i> (aspen) <i>Populus balsamifera</i> (balsam poplar) <i>Lonicera dioica</i> (twining honeysuckle) <i>Abies balsamea</i> (balsam fir) <i>Abies lasiocarpa</i> (subalpine fir) <i>Lonicera involucrata</i> (bracted honeysuckle)

Bracted honeysuckle

This ecosite is characterized by subhygric to mesic moisture regimes and rich nutrient levels. Indicator species include balsam poplar, bracted honeysuckle, dogwood, currant, devil's-club, wild red raspberry, tall lungwort, sweet-scented bedstraw, oak fern, lady fern, and narrow spinulose shield fern. This ecosite has four ecosite phases, (f1) bracted honeysuckle PI, (f2) bracted honeysuckle Aw-Pb, (f3) bracted honeysuckle Aw-Sw-PI, and (f4) bracted honeysuckle Sw.

(f) Bracted honeysuckle – characteristic woody plant species.

Tree	<i>Pinus contorta</i> (lodgepole pine) <i>Abies balsamea</i> (balsam fir) <i>Abies lasiocarpa</i> (subalpine fir) <i>Picea glauca</i> (white spruce) <i>Picea mariana</i> (black spruce) <i>Populus tremuloides</i> (aspen) <i>Betula papyrifera</i> (white birch) <i>Populus balsamifera</i> (balsam poplar)
Shrub	<i>Alnus crispa</i> (green alder) <i>Abies balsamea</i> (balsam fir) <i>Abies lasiocarpa</i> (subalpine fir) <i>Lonicera involucrata</i> (bracted honeysuckle) <i>Oplopanax horridum</i> (devil's-club)



Viburnum edule (low-bush cranberry)
Rosa acicularis (prickly rose)
Ledum groenlandicum (Labrador tea)
Rubus idaeus (wild red raspberry)
Linnaea borealis (twin-flower)
Vaccinium membranaceum (tall bilberry)
Rubus parviflorus (thimbleberry)
Cornus stolonifera (dogwood)
Populus tremuloides (aspen)
Populus balsamifera (balsam poplar)
Ribes spp. (currant)
Betula papyrifera (white birch)
Salix spp. (willow)
Picea glauca (white spruce)
Rubus parviflorus (thimbleberry)
Spiraea betulifolia (white meadowsweet)
Alnus tenuifolia (river alder)

Meadow

This ecosite is characterized by subhydryc to mesic moisture regimes and rich to very rich nutrient levels. Indicator species include willow, cow parsnip, veiny meadow rue, tall lungwort, long-stalked chickweed, avens, marsh reed grass and fringed brome. This ecosite has two ecosite phases, (g1) shrubby meadow and (g2) forb meadow. There are no characteristic woody plant species found in (g2) forb meadow.

(g) Meadow – characteristic woody plant species (g1 only).

Tree	No tree species
Shrub	<i>Salix</i> spp. (willow) <i>Betula pumila</i> (dwarf birch) <i>Ribes</i> spp. (currant) <i>Alnus tenuifolia</i> (river alder)

Labrador tea–subhygric

This ecosite is characterized by hygric to subhygric moisture regimes and poor to very poor nutrient levels. Indicator species include lodgepole pine, black spruce, Labrador tea and bog cranberry. This ecosite has one ecosite phase, (h1) Labrador tea–subhygric Sb-Pl.

(h) Labrador tea–subhygric – characteristic woody plant species.

Tree	<i>Pinus contorta</i> (lodgepole pine) <i>Picea mariana</i> (black spruce) <i>Picea glauca</i> (white spruce)
Shrub	<i>Ledum groenlandicum</i> (Labrador tea)



Alnus crispa (green alder)
Vaccinium vitis-idaea (bog cranberry)
Picea mariana (black spruce)
Vaccinium myrtilloides (blueberry)

Horsetail

This ecosite is characterized by hygric to subhygric moisture regimes and rich to medium nutrient levels. Indicator species include common horsetail and meadow horsetail. This ecosite has three ecosite phases, (i1) horsetail Pb-Aw, (i2) horsetail Pb-Sw, and (i3) horsetail Sw.

(i) Horsetail – characteristic woody plant species.

Tree	<i>Populus tremuloides</i> (aspen) <i>Populus balsamifera</i> (balsam poplar) <i>Picea glauca</i> (white spruce) <i>Betula papyrifera</i> (white birch) <i>Abies balsamea</i> (balsam fir) <i>Abies lasiocarpa</i> (subalpine fir)
Shrub	<i>Rosa acicularis</i> (prickly rose) <i>Rubus idaeus</i> (wild red raspberry) <i>Salix</i> spp. (willow) <i>Amelanchier alnifolia</i> (saskatoon) <i>Populus tremuloides</i> (aspen) <i>Populus balsamifera</i> (balsam poplar) <i>Symphoricarpos albus</i> (snowberry) <i>Ribes</i> spp. (currant) <i>Linnaea borealis</i> (twin-flower) <i>Viburnum edule</i> (low-bush cranberry) <i>Betula papyrifera</i> (white birch) <i>Picea glauca</i> (white spruce) <i>Shepherdia canadensis</i> (Canada buffalo-berry) <i>Lonicera involucrata</i> (bracted honeysuckle)

Labrador tea/horsetail

This ecosite is characterized by subhydryc to hygric moisture regimes and rich to poor nutrient levels. Indicator species include black spruce, white spruce, Labrador tea, common horsetail, woodland horsetail and meadow horsetail. This ecosite has one ecosite phase, (j1) Labrador tea/horsetail Sb-Sw.

(j) Labrador tea/horsetail – characteristic woody plant species.

Tree	<i>Picea mariana</i> (black spruce) <i>Picea glauca</i> (white spruce)
Shrub	<i>Ledum groenlandicum</i> (Labrador tea)



Lonicera involucrata (bracted honeysuckle)
Rosa acicularis (prickly rose)
Vaccinium vitis-idaea (bog cranberry)

Bog

This ecosite is characterized by hydric to hygric moisture regimes and poor to very poor nutrient levels. Indicator species include black spruce, Labrador tea, bog cranberry, small bog cranberry, cloudberry and peat moss. This ecosite has two ecosite phases, (k1) treed bog, and (k2) shrubby bog.

(k) Bog – characteristic woody plant species.

Tree	<i>Picea mariana</i> (black spruce)
Shrub	<i>Ledum groenlandicum</i> (Labrador tea) <i>Picea mariana</i> (black spruce) <i>Salix</i> spp. (willow) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Oxycoccus microcarpus</i> (small bog cranberry) <i>Vaccinium myrtilloides</i> (blueberry) <i>Pinus contorta</i> (lodgepole pine) <i>Betula pumila</i> (dwarf birch)

Poor fen

This ecosite is characterized by hydric to hygric moisture regimes and rich to poor nutrient levels. Indicator species include black spruce, tamarack, Labrador tea, dwarf birch, willow, cloudberry, sedge, peat moss, golden moss, tufted moss and brown moss. This ecosite has two ecosite phases, (l1) treed poor fen and (l2) shrubby poor fen.

(l) Poor fen – characteristic woody plant species.

Tree	<i>Picea mariana</i> (black spruce) <i>Larix laricina</i> (tamarack)
Shrub	<i>Ledum groenlandicum</i> (Labrador tea) <i>Betula pumila</i> (dwarf birch) <i>Picea mariana</i> (black spruce) <i>Salix</i> spp. (willow) <i>Larix laricina</i> (tamarack) <i>Vaccinium vitis-idaea</i> (bog cranberry) <i>Oxycoccus microcarpus</i> (small bog cranberry)



Rich fen

This ecosite is characterized by hydric to hygric moisture regimes and medium to very rich nutrient levels. Indicator species include tamarack, willow, dwarf birch, sedge, golden moss, tufted moss, brown moss, buck-bean and marsh marigold. This ecosite has three ecosite phases, (m1) treed rich fen, (m2) shrubby rich fen, and (m3) graminoid rich fen.

(m) Rich fen – characteristic woody plant species.

Tree	<i>Larix laricina</i> (tamarack) <i>Picea mariana</i> (black spruce)
Shrub	<i>Betula pumila</i> (dwarf birch) <i>Ledum groenlandicum</i> (Labrador tea) <i>Salix</i> spp. (willow) <i>Larix laricina</i> (tamarack) <i>Ribes</i> spp. (currant) <i>Picea mariana</i> (black spruce) <i>Salix</i> spp. (willow)

Marsh

This ecosite is characterized by hydric to subhydric moisture regimes and rich to very rich nutrient levels. Indicator species include common cattail, rush, sedge and bulrush. This ecosite has one ecosite phase, (n1) marsh. This ecosite is not characterized by any woody species.

3. Autecology of key Boreal Mixedwood species

Adapted from *Alberta Silviculture Guide* (2007)²

² Geographic Dynamics Corp. and Incremental Forest Technologies Ltd., 2007. ***Alberta Silviculture Guide: Boreal Mixedwood and Lower Foothills Natural Subregions. Version 1.*** Developed by Geographic Dynamics Corp. and Incremental Forest Technologies with contributions from Coast to Coast Reforestation Inc. Developed for the Mixedwood Management Association, Edmonton, Alberta.



Target Plant Communities

Autecological Factor	White spruce	Black spruce	Aspen	Balsam poplar	Alder	Willow	Labrador Tea	Beaked Hazel
Sere	Mid	Plastic	Pioneer	Pioneer	Pioneer	Pioneer to late	Plastic	Pioneer to late
Tolerance	Mid.	Int.	Int.	Int.	Int.	Varies with species	Tol.	Int.
Reproduction	Seed	Seed	Suckers	Sprouts	Seed	Seed	Seed, Sprouts	Seed, Suckers, Sprouting
		Layer	Sprouts	Seeds	Sprouts	Seed bank		
			Seed	Suckers		Sprouts		
				Cuttings				
Optimal moisture regime	Mesic to sub-hygric	Mesic to sub-hydric	Mesic	Moist to sub-hygric	Submesic to hygric	Variable	Hygric	Mesic to hygric
Drought tolerance	Medium	Low	Low	Low	High	Variable	Medium	Medium
Site richness	Medium to rich	Low to medium	Medium to rich	Rich to medium	Medium to rich	Medium to rich	Low to Medium	Medium to rich



Target Plant Communities

Autecological Factor	White spruce	Black spruce	Aspen	Balsam poplar	Alder	Willow	Labrador Tea	Beaked Hazel
Response to:								
Wildfire	Loss of site	Invasion	Inundatory invasion	Invasion by seed	Inundatory invasion	Loss of site	Increases in abundance	Variable depending on root damage
Ground fire	Invasion	Site dominance	Loss of site or invasion	Loss of site	Re-invasion	Loss of site	Invades from seed	Loss of site
							Sprouts	Stimulates sprouting
Wildlife habitat value			Birds and mammals: breeding, foraging, feeding -Moose (Alces alces)					



BOREAL RESEARCH INSTITUTE

Target Plant Communities